

# ENDOMORPHISM ALGEBRAS OF 2-TERM SILTING COMPLEXES

ASLAK BAKKE BUAN AND YU ZHOU

**ABSTRACT.** We study possible values of the global dimension of endomorphism algebras of 2-term silting complexes. We show that for any algebra  $A$  whose global dimension  $\text{gl.dim } A \leq 2$  and any 2-term silting complex  $\mathbf{P}$  in the bounded derived category  $D^b(A)$  of  $A$ , the global dimension of  $\text{End}_{D^b(A)}(\mathbf{P})$  is at most 7. We also show that for each  $n > 2$ , there is an algebra  $A$  with  $\text{gl.dim } A = n$  such that  $D^b(A)$  admits a 2-term silting complex  $\mathbf{P}$  with  $\text{gl.dim } \text{End}_{D^b(A)}(\mathbf{P})$  infinite.

## INTRODUCTION

Let  $A$  be a finite dimensional algebra over a field  $k$ . Let  $T$  be a (classical) tilting module in the category  $\text{mod } A$  of finite dimensional right  $A$ -modules; that is the projective dimension  $\text{pd } T$  is at most 1, we have  $\text{Ext}_A^1(T, T) = 0$  and there is an exact sequence  $0 \rightarrow A \rightarrow T_1 \rightarrow T_2 \rightarrow 0$  with  $T_1, T_2$  in  $\text{add } T$ , the additive closure of  $T$ . Let  $B = \text{End}_A(T)$ . Then, it is a well-known fact (see for example [8, III, Section 3.4] for a more general statement) that  $\text{gl.dim } B \leq \text{gl.dim } A + 1$ , where  $\text{gl.dim } A$  denotes the global dimension of  $A$ .

In this paper we investigate to which extent this generalizes to the following setting. We now consider a 2-term silting complex  $\mathbf{P}$  in the bounded homotopy category of finitely generated projective  $A$ -modules,  $K^b(\text{proj } A)$ . This is just a map between projective  $A$ -modules, considered as a complex, with the property that  $\text{Hom}_{K^b(\text{proj } A)}(\mathbf{P}, \mathbf{P}[1]) = 0$  where  $[1]$  denotes the shift functor, and such that  $\mathbf{P}$  generates  $K^b(\text{proj } A)$  as a triangulated category. Note that  $K^b(\text{proj } A)$  can be considered to be a full triangulated subcategory of the derived category  $D^b(A)$ .

The concept of silting complexes originated from [11], and has more recently been studied by many authors, often motivated by combinatorial aspects related to mutations, as in [2]. Moreover, the case of 2-term silting is of particular interest, see e.g. [1], [4] and [12].

In the setting of 2-term silting, we have the following theorem:

**Theorem 0.1.** *Let  $B = \text{End}_{D^b(A)}(\mathbf{P})$ , for a 2-term silting complex  $\mathbf{P}$  in  $K^b(\text{proj } A)$ . Then the following hold.*

- (a) *If  $\text{gl.dim } A = 1$ , then  $\text{gl.dim } B \leq 3$ .*
- (b) *If  $\text{gl.dim } A = 2$ , then  $\text{gl.dim } B \leq 7$ .*

*Moreover, for each  $n > 2$ , there is an algebra  $A$ , with  $\text{gl.dim } A = n$ , such that  $K^b(\text{proj } A)$  admits a 2-term silting complex  $\mathbf{P}$  with  $\text{gl.dim } \text{End}_{D^b(A)}(\mathbf{P}) = \infty$ .*

Note that the projective presentation of a tilting  $A$ -module  $T$  as defined above, gives rise to a 2-term silting complex  $\mathbf{P}_T$  in  $K^b(\text{proj } A)$ , and that we have an isomorphism of algebras  $\text{End}_A(T) \cong \text{End}_{D^b(A)}(\mathbf{P}_T)$ .

The situation in part (a) was studied in [6]. In this case  $B$  is called a *silted algebra*, and it was proved that silted algebras are so-called shod algebras [7], in particular this implies that  $\text{gl.dim } B \leq 3$ , by [9].

---

This work was supported by FRINAT grant number 231000, from the Norwegian Research Council. Support by the Institut Mittag-Leffler (Djursholm, Sweden) is gratefully acknowledged.

The main body of this paper is a proof of (b), an example that the global dimension of  $B$  actually can be 7 in this case, and a class of examples that justifies the last statement of Theorem 0.1.

We also prove that with a stronger assumption on  $\mathbf{P}$ , we actually get that  $\text{gl.dim } B$  is bounded by  $\text{gl.dim } A$ . More precisely, we show the following.

**Theorem 0.2.** *With the above notation, and assuming in addition that  $\text{pd } H^0(\mathbf{P}) \leq 1$ , we have  $\text{gl.dim } B \leq 2(\text{gl.dim } A) + 2$ .*

In the first section, we recall some notation and background concerning 2-term silting complexes and their endomorphism algebras. In the second section, we prove some preliminary general results. Then, in Section 3 and 4, we prove respectively Theorem 0.2 and Theorem 0.1, while in the last section, we give some examples.

## 1. BACKGROUND AND NOTATION

Let  $A$  be a finite dimensional algebra with  $\text{gl.dim } A = d$ . Then  $K^b(\text{proj } A) = D^b(A) := \mathcal{D}$ . Let  $\mathbf{P}$  be a 2-term silting complex in  $\mathcal{D}$  and let  $B = \text{End}_{\mathcal{D}}(\mathbf{P})$ . We recall some classical notation (see e.g. [3]) and some results from [5], which will be used freely in the remaining of the paper.

Recall that a pair of subcategories  $(\mathcal{X}, \mathcal{Y})$  of  $\text{mod } A$ , is called a *torsion pair*, if the following hold:

- $\text{Hom}_A(\mathcal{X}, Y) = 0$  if and only if  $Y$  is in  $\mathcal{Y}$ , and
- $\text{Hom}_A(X, \mathcal{Y}) = 0$  if and only if  $X$  is in  $\mathcal{X}$ .

For a given torsion pair  $(\mathcal{X}, \mathcal{Y})$  and an object  $M$  in  $\text{mod } A$ , there is a (unique) exact sequence

$$0 \rightarrow tM \rightarrow M \rightarrow M/tM \rightarrow 0$$

with  $tM$  in  $\mathcal{X}$  and  $M/tM$  in  $\mathcal{Y}$ . This is called the *canonical sequence* of  $M$ . Furthermore, for an  $A$ -module  $X$  we let  $\text{add } X$  denote the additive closure of  $X$  in  $\text{mod } A$ , and we let  $\text{Fac } X$  denote the full subcategory of all quotients of modules in  $\text{add } X$ . The first notion is also used for a complex  $X$  in  $\mathcal{D}$ .

For a 2-term silting complex  $\mathbf{P}$ , consider the full subcategories of  $\text{mod } A$  given by

- $\mathcal{T}(\mathbf{P}) = \{X \in \text{mod } A \mid \text{Hom}_{\mathcal{D}}(\mathbf{P}, X[1]) = 0\}$ , and
- $\mathcal{F}(\mathbf{P}) = \{Y \in \text{mod } A \mid \text{Hom}_{\mathcal{D}}(\mathbf{P}, Y) = 0\}$ .

Furthermore, let  $B = \text{End}_{\mathcal{D}}(\mathbf{P})$ . The following summarizes results from [5] which will be essential later in this paper.

**Proposition 1.1.** *Let  $\mathbf{P}$  be a 2-term silting complex in  $K^b(\text{proj } A)$ . Then the following hold.*

- (a) *The pair  $(\mathcal{T}(\mathbf{P}), \mathcal{F}(\mathbf{P}))$  is a torsion pair in  $\text{mod } A$ .*
- (b)  *$\mathcal{T}(\mathbf{P}) = \text{Fac } H^0(\mathbf{P})$ .*
- (c) *The category  $C(\mathbf{P}) = \{X \in \mathcal{D} \mid \text{Hom}(\mathbf{P}, X[i]) = 0 \text{ for } i \neq 0\}$  is an abelian category with short exact sequences coinciding with the triangles in  $\mathcal{D}$  whose vertices are in  $C(\mathbf{P})$ .*
- (d) *Let  $X$  be in  $\mathcal{D}$ . Then we have that  $X$  is in  $C(\mathbf{P})$  if and only if  $H^0(X)$  is in  $\mathcal{T}(\mathbf{P})$ ,  $H^{-1}(X)$  is in  $\mathcal{F}(\mathbf{P})$  and  $H^i(X) = 0$  for  $i \neq -1, 0$ .*
- (e)  *$\text{Hom}_{\mathcal{D}}(\mathbf{P}, -): C(\mathbf{P}) \rightarrow \text{mod } B$  is an equivalence of (abelian) categories.*

For full subcategories  $\mathcal{X}$  and  $\mathcal{Y}$  of  $\mathcal{D}$ , we let  $\mathcal{X} * \mathcal{Y}$  denote the full subcategory of  $\mathcal{D}$  with objects  $Z$  appearing in a triangle

$$X \rightarrow Z \rightarrow Y \rightarrow X[1]$$

with  $X$  in  $\mathcal{X}$  and  $Y$  in  $\mathcal{Y}$ . It follows from the octahedral axiom that we have  $(\mathcal{X} * \mathcal{Y}) * \mathcal{Z} = \mathcal{X} * (\mathcal{Y} * \mathcal{Z})$ , for three full subcategories  $\mathcal{X}, \mathcal{Y}$  and  $\mathcal{Z}$ . The subcategory  $\mathcal{X}$  is called *extension*

closed if  $\mathcal{X} * \mathcal{X} = \mathcal{X}$ . We will need the following fact, which follows from [10, Propositions 2.1 and 2.4].

**Lemma 1.2.** *Let  $\mathcal{X}_i$  be subcategories of  $\mathcal{D}$ , with  $\text{Hom}_{\mathcal{D}}(\mathcal{X}_i, \mathcal{X}_j) = 0 = \text{Hom}_{\mathcal{D}}(\mathcal{X}_i, \mathcal{X}_j[1])$  for  $i < j$ . Then  $\mathcal{X}_1 * \mathcal{X}_2 * \cdots * \mathcal{X}_n$  is closed under extensions and direct summands.*

## 2. PRELIMINARIES

Now, fix a 2-term sifting complex  $\mathbf{P}$  in  $K^b(\text{proj } A)$ , and let  $\mathcal{P} = \text{add } \mathbf{P}$ . In this section we include some general observations on projective objects and projective dimensions in  $C(\mathbf{P})$ .

For each  $\mathbf{P}_0$  in  $\mathcal{P}$ , given by  $P_0^{-1} \xrightarrow{p_0} P_0^0$ , consider the canonical exact sequence of  $H^{-1}(\mathbf{P}_0)$  relative to the torsion pair  $(\mathcal{T}(\mathbf{P}), \mathcal{F}(\mathbf{P}))$ :

$$0 \rightarrow tH^{-1}(\mathbf{P}_0) \rightarrow H^{-1}(\mathbf{P}_0) \rightarrow H^{-1}(\mathbf{P}_0)/tH^{-1}(\mathbf{P}_0) \rightarrow 0.$$

So  $tH^{-1}(\mathbf{P}_0)$  is a submodule of  $P_0^{-1}$  and we denote by  $\pi: P_0^{-1} \rightarrow P_0^{-1}/tH^{-1}(\mathbf{P}_0)$  the canonical epimorphism. Let  $\tilde{\mathbf{P}}_0$  be the complex  $P_0^{-1}/tH^{-1}(\mathbf{P}_0) \xrightarrow{\tilde{p}_0} P_0^0$ , where  $\tilde{p}_0$  is the unique homomorphism such that the diagram

$$\begin{array}{ccc} & P_0^{-1}/tH^{-1}(\mathbf{P}_0) & \\ \pi \nearrow & & \searrow \tilde{p}_0 \\ P_0^{-1} & \xrightarrow{p_0} & P_0^0 \end{array}$$

commutes.

Let  $\mathcal{P}_C = \mathcal{P} \cap C(\mathbf{P})$ .

**Lemma 2.1.** *Let  $\mathbf{P}_0$  be in  $\mathcal{P}$ . Then  $\mathbf{P}_0$  is in  $\mathcal{P}_C$  if and only if  $\mathbf{P}_0 \cong \tilde{\mathbf{P}}_0$ .*

*Proof.* We have by definition that  $\mathbf{P}_0 \cong \tilde{\mathbf{P}}_0$  if and only if  $tH^{-1}(\mathbf{P}_0) = 0$  if and only if  $H^{-1}(\mathbf{P}_0)$  is in  $\mathcal{F}(\mathbf{P})$  if and only if  $\text{Hom}(\mathbf{P}, H^{-1}(\mathbf{P}_0)) = 0$ . It is straightforward to check that  $\text{Hom}(\mathbf{P}, H^{-1}(\mathbf{P}_0)) = 0$  if and only if  $\text{Hom}(\mathbf{P}, \mathbf{P}_0[-1]) = 0$ . Moreover, we have that  $\text{Hom}(\mathbf{P}, \mathbf{P}_0[-1]) = 0$  if and only if  $\mathbf{P}_0$  is in  $C(\mathbf{P})$ , and the statement follows from this.  $\square$

**Lemma 2.2.** *With notation as above, the following hold.*

(a) *There is a triangle in  $\mathcal{D}$ :*

$$tH^{-1}(\mathbf{P})[1] \rightarrow \mathbf{P} \rightarrow \tilde{\mathbf{P}} \rightarrow tH^{-1}(\mathbf{P})[2].$$

(b) *The object  $\tilde{\mathbf{P}}$  is a projective generator for  $C(\mathbf{P})$ .*

*Proof.* The triangle in (a) exists by the construction of  $\tilde{\mathbf{P}}$ .

Note that  $H^0(\tilde{\mathbf{P}}) = H^0(\mathbf{P})$  is in  $\mathcal{T}(\mathbf{P})$  and  $H^{-1}(\tilde{\mathbf{P}}) = H^{-1}(\mathbf{P})/tH^{-1}(\mathbf{P})$  is in  $\mathcal{F}(\mathbf{P})$ . Then by Proposition 1.1 (d), we have  $\tilde{\mathbf{P}} \in C(\mathbf{P})$ . Applying the functor  $\text{Hom}_{\mathcal{D}}(\mathbf{P}, -)$  to this triangle yields an isomorphism

$$\text{Hom}_{\mathcal{D}}(\mathbf{P}, \mathbf{P}) \cong \text{Hom}_{\mathcal{D}}(\mathbf{P}, \tilde{\mathbf{P}})$$

as  $B$ -modules. Now (b) follows from Proposition 1.1 (e).  $\square$

For any integer  $i$ , we let  $\mathcal{D}^{\leq i}(\mathbf{P}) = \{\mathbf{X} \in \mathcal{D} \mid \text{Hom}_{\mathcal{D}}(\mathbf{P}, \mathbf{X}[j]) = 0 \text{ for } j > i\}$ , and we let  $\mathcal{D}^{\geq i}(\mathbf{P}) = \{\mathbf{X} \in \mathcal{D} \mid \text{Hom}_{\mathcal{D}}(\mathbf{P}, \mathbf{X}[j]) = 0 \text{ for } j < i\}$ .

**Lemma 2.3.** *With notation as above, we have:  $C(\mathbf{P}) \subset \mathcal{P} * \mathcal{P}[1] * \cdots * \mathcal{P}[d+1]$ .*

*Proof.* By [2, Proposition 2.23], we have

$$C(\mathbf{P}) \subset \mathcal{D}^{\leq 0}(\mathbf{P}) \subset \mathcal{P} * \mathcal{P}[1] * \cdots * \mathcal{P}[l-1] * \mathcal{P}[l]$$

for some  $l > 0$ . For any  $\mathbf{M}$  in  $C(\mathbf{P})$ , by Proposition 1.1 (d), we have  $H^i(\mathbf{M}) = 0$  for  $i \neq -1, 0$ . So there is a complex  $\mathbf{X}$  of projective  $A$ -modules, which is equivalent to  $\mathbf{M}$ , and such that  $H^i(\mathbf{X}) = 0$  for  $i > 0$  or  $i < -d - 1$ . So

$$\mathrm{Hom}_{\mathcal{D}}(\mathbf{M}, \mathbf{P}[i]) \cong \mathrm{Hom}_{\mathcal{D}}(\mathbf{X}, \mathbf{P}[i]) = 0, \quad i \geq d + 2,$$

which implies that  $\mathbf{M}$  is in  $\mathcal{P} * \mathcal{P}[1] * \cdots * \mathcal{P}[d + 1]$ .  $\square$

**Lemma 2.4.** *For a complex  $\mathbf{X}$  in  $C(\mathbf{P}) \cap (\mathcal{P}_C * \mathcal{P}_C[1] * \cdots * \mathcal{P}_C[m])$  for some  $m \geq 0$ , we have  $\mathrm{pd} \mathrm{Hom}_{\mathcal{D}}(\mathbf{P}, \mathbf{X})_B \leq m$ .*

*Proof.* Let  $\mathbf{X}_0 = \mathbf{X}$ . There are triangles

$$\mathbf{X}_{i+1} \rightarrow \mathbf{O}_i \xrightarrow{g_i} \mathbf{X}_i \rightarrow \mathbf{X}_{i+1}[1], \quad 0 \leq i \leq m - 1$$

where  $\mathbf{O}_i$  is in  $\mathcal{P}_C$  and  $\mathbf{X}_i$  is in  $\mathcal{P}_C * \mathcal{P}_C[1] * \cdots * \mathcal{P}_C[m - i]$ . Since  $\mathrm{Hom}_{\mathcal{D}}(\mathbf{P}, \mathbf{P}[i]) = 0$  for all  $i > 0$ , we have that  $g_i$  is a right  $\mathcal{P}$ -approximation of  $\mathbf{X}_i$ . By Lemma 2.1 and Lemma 2.2 (b), each  $\mathbf{O}_i$  is projective in  $C(\mathbf{P})$ . Assume that  $\mathbf{X}_i$  is in  $C(\mathbf{P})$  for some  $0 \leq i \leq m - 1$ . Then, since  $g_i$  is a right  $\mathcal{P}$ -approximation and  $\mathbf{O}_i$  is projective in  $C(\mathbf{P})$ , we have that  $g_i$  is an epimorphism in  $C(\mathbf{P})$ . So  $\mathbf{X}_{i+1}$  is the kernel of  $g_i$ , by Proposition 1.1 (c). Note that  $\mathbf{X}_0 \in C(\mathbf{P})$ . Then by induction on  $i$ , we have that  $\mathbf{X}_i \in C(\mathbf{P})$  for all  $0 \leq i \leq m$  and

$$\mathrm{pd} \mathrm{Hom}_{\mathcal{D}}(\mathbf{P}, \mathbf{X}_i)_B \leq \mathrm{pd} \mathrm{Hom}_{\mathcal{D}}(\mathbf{P}, \mathbf{X}_{i+1})_B + 1.$$

Therefore  $\mathrm{pd} \mathrm{Hom}_{\mathcal{D}}(\mathbf{P}, \mathbf{X})_B \leq \mathrm{pd} \mathrm{Hom}_{\mathcal{D}}(\mathbf{P}, \mathbf{X}_m)_B + m = m$  since  $\mathbf{X}_m \in \mathcal{P}_C$  is projective in  $C(\mathbf{P})$ .  $\square$

We end this section by considering the following special case. Recall from [13], that a 2-term silting complex  $\mathbf{P}$  in  $K^b(\mathrm{proj} A)$  is a *tilting complex* if  $\mathrm{Hom}_{\mathcal{D}}(\mathbf{P}, \mathbf{P}[-1]) = 0$ .

**Proposition 2.5.** *If the 2-term silting complex  $\mathbf{P}$  is a tilting complex, then  $\mathrm{gl.dim} \mathrm{End}_{\mathcal{D}}(\mathbf{P}) \leq \mathrm{gl.dim} A + 1$ .*

*Proof.* If  $\mathbf{P}$  is tilting, then  $\mathbf{P}$  is in  $C(\mathbf{P})$ . So we infer that  $\mathcal{P} = \mathcal{P}_C$ . It follows from Lemma 2.3 and Lemma 2.4 that  $\mathrm{gl.dim} \mathrm{End}_{\mathcal{D}}(\mathbf{P}) \leq \mathrm{gl.dim} A + 1$ .  $\square$

Note that the classical situation (as in [8, III, section 3.4]) where  $\mathbf{P}$  is the projective resolution of a classical tilting module, is covered by this result.

### 3. THE PARTIAL TILTING CASE

Throughout this section, we assume that  $\mathrm{pd} H^0(\mathbf{P})_A \leq 1$ , that is:  $H^0(\mathbf{P})$  is a partial tilting  $A$ -module. Then we have that  $Q = H^{-1}(\mathbf{P})$  is projective as an  $A$ -module, and  $\mathbf{P} \cong H^0(\mathbf{P}) \oplus Q[1]$ . Consider the canonical exact sequence of  $Q$  relative to the torsion pair  $(\mathcal{T}(\mathbf{P}), \mathcal{F}(\mathbf{P}))$ :

$$0 \rightarrow tQ \rightarrow Q \rightarrow Q/tQ \rightarrow 0.$$

As before, we let  $d = \mathrm{gl.dim} A$ . We first prove two technical lemmas.

**Lemma 3.1.** *With the above notation, we have*

$$tQ \in \mathrm{add} H^0(\mathbf{P}) * \mathrm{add} H^0(\mathbf{P})[1] * \cdots * \mathrm{add} H^0(\mathbf{P})[d - 1].$$

*Proof.* We first note that  $tQ \in \mathcal{T}(\mathbf{P})$ , so by definition  $\mathrm{Hom}_{D^b(A)}(\mathbf{P}, tQ[i]) = 0$  for  $i \neq 0$ . In particular, we have  $\mathrm{Hom}_{D^b(A)}(Q[1], tQ[i]) = 0$  for  $i \neq 0$ . For  $i = 0$ , since both  $Q$  and  $tQ$  are  $A$ -modules, we also have that  $\mathrm{Hom}_{D^b(A)}(Q[1], tQ) = 0$ . It follows from  $T(\mathbf{P}) \in C(\mathbf{P})$  that by Proposition 1.1, we have  $tQ \in \mathcal{P} * \mathcal{P}[1] * \cdots * \mathcal{P}[d+1]$ . Therefore, using  $\mathbf{P} \cong H^0(\mathbf{P}) \oplus Q[1]$ , we get that  $tQ$  is in  $\mathrm{add} H^0(\mathbf{P}) * \mathrm{add} H^0(\mathbf{P})[1] * \cdots * \mathrm{add} H^0(\mathbf{P})[d+1]$ . By the canonical sequence of  $Q$ , we have  $\mathrm{pd}(tQ)_A \leq \mathrm{pd}(Q/tQ)_A - 1 \leq d - 1$ . Hence, it follows that  $\mathrm{Hom}(tQ, \mathbf{P}[d]) = 0 = \mathrm{Hom}(tQ, \mathbf{P}[d+1])$ . The claim of the lemma follows.  $\square$

**Lemma 3.2.** *With the above notation, we have  $C(\mathbf{P}) \subset \mathcal{P} * \mathcal{P}[1] * \cdots * \mathcal{P}[d] * \mathrm{add} H^0(\mathbf{P})[d+1]$ .*

*Proof.* Using that  $\mathbf{P} \cong H^0(\mathbf{P}) \oplus Q[1]$  in combination with Lemma 2.3, we only need to prove that  $\mathrm{Hom}_{\mathcal{D}}(\mathbf{X}, Q[d+2]) = 0$  for  $\mathbf{X} \in C(\mathbf{P})$ . This follows from  $\mathrm{pd} H^i(\mathbf{X})_A \leq d$  for  $i = -1, 0$  and  $H^i(\mathbf{X}) = 0$  for  $i \neq -1, 0$ .  $\square$

We can now prove the main result of this section.

**Theorem 3.3.** *If  $\mathrm{pd}(H^0(\mathbf{P}))_A \leq 1$ , then  $\mathrm{gl.dim} B \leq 2 \mathrm{gl.dim} A + 2$ .*

*Proof.* Let  $\mathbf{X}$  be an object in  $C(\mathbf{P})$  with

$$\mathbf{X} \in \mathcal{P} * \cdots * \mathcal{P}[i] * \mathrm{add} H^0(\mathbf{P})[i+1] * \cdots * \mathrm{add} H^0(\mathbf{P})[d+1]$$

for some  $0 \leq i \leq d$ . Then there is a triangle

$$\mathbf{X}_1 \rightarrow \mathbf{E} \xrightarrow{g_{\mathbf{X}}} \mathbf{X} \rightarrow \mathbf{X}_1[1]$$

where  $g_{\mathbf{X}}$  is a right  $\mathcal{P}$ -approximation of  $\mathbf{X}$  and  $\mathbf{X}_1$  is in

$$\mathcal{P} * \cdots * \mathcal{P}[i-1] * \mathrm{add} H^0(\mathbf{P})[i] * \cdots * \mathrm{add} H^0(\mathbf{P})[d].$$

Then  $\mathrm{Hom}_{\mathcal{D}}(\mathbf{P}, g_{\mathbf{X}})$  is an epimorphism and  $\mathrm{Hom}_{\mathcal{D}}(\mathbf{P}, \mathbf{E})$  is projective in  $\mathrm{mod} B$ .

Recall that  $Q = H^{-1}(\mathbf{P})$ . Then, by Lemma 2.2 there is a triangle

$$F[1] \rightarrow \mathbf{E} \rightarrow \widetilde{\mathbf{E}} \rightarrow F[2]$$

where  $F$  is in  $\mathrm{add} tQ \subset \mathcal{T}(\mathbf{P}) \subset C(\mathbf{P})$  and  $\widetilde{\mathbf{E}}$  is projective in  $C(\mathbf{P})$ . So  $\mathrm{Hom}_{\mathcal{D}}(F[1], \mathbf{X}) = 0$  since  $\mathbf{X} \in C(\mathbf{P})$ . It follows that the map  $g_{\mathbf{X}}$  factors through the map  $\mathbf{E} \rightarrow \widetilde{\mathbf{E}}$ . Then, by the octahedral axiom, we have the following commutative diagram of triangles:

$$\begin{array}{ccccccc} & & \mathbf{X}[-1] & \xlongequal{\quad} & \mathbf{X}[-1] & & \\ & & \downarrow & & \downarrow & & \\ F[1] & \longrightarrow & \mathbf{X}_1 & \longrightarrow & \mathbf{X}' & \longrightarrow & F[2] \\ \parallel & & \downarrow & & \downarrow & & \parallel \\ F[1] & \longrightarrow & \mathbf{E} & \longrightarrow & \widetilde{\mathbf{E}} & \longrightarrow & F[2] \\ & & \downarrow g_{\mathbf{X}} & & \downarrow \widetilde{g}_{\mathbf{X}} & & \\ & & \mathbf{X} & \xlongequal{\quad} & \mathbf{X} & & \end{array}.$$

Then we have that

$$\begin{aligned} \mathbf{X}' &\in \mathrm{add} \mathbf{X}_1 * \mathrm{add} F[2] \\ &\subset (\mathcal{P} * \cdots * \mathcal{P}[i-1] * \mathrm{add} H^0(\mathbf{P})[i] * \cdots * \mathrm{add} H^0(\mathbf{P})[d]) \\ &\quad * (\mathrm{add} H^0(\mathbf{P}) * \cdots * \mathrm{add} H^0(\mathbf{P})[d-1])[2] \\ &= (\mathcal{P} * \cdots * \mathcal{P}[i-1] * \mathrm{add} H^0(\mathbf{P})[i] * \cdots * \mathrm{add} H^0(\mathbf{P})[d]) * \mathrm{add} H^0(\mathbf{P})[d+1] \end{aligned}$$

where the inclusion is due to Lemma 3.1, and the equality follows from

$$\mathcal{P} * \cdots * \mathcal{P}[i-1] * \text{add } H^0(\mathbf{P})[i] * \cdots * \text{add } H^0(\mathbf{P})[d]$$

being closed under extensions by Lemma 1.2. Applying  $\text{Hom}_{\mathcal{D}}(\mathbf{P}, -)$  to the above diagram, we obtain a commutative diagram

$$\begin{array}{ccc} \text{Hom}_{\mathcal{D}}(\mathbf{P}, \mathbf{E}) & \xrightarrow{\cong} & \text{Hom}_{\mathcal{D}}(\mathbf{P}, \widetilde{\mathbf{E}}) \\ & \searrow \text{Hom}_{\mathcal{D}}(\mathbf{P}, g_X) & \swarrow \text{Hom}_{\mathcal{D}}(\mathbf{P}, \widetilde{g}_X) \\ & \text{Hom}_{\mathcal{D}}(\mathbf{P}, \mathbf{X}) & \end{array}$$

Using that the map  $\text{Hom}_{\mathcal{D}}(\mathbf{P}, g_X)$  is an epimorphism in  $\text{mod } B$ , it follows that the map  $\widetilde{g}_X$  is an epimorphism in  $C(\mathbf{P})$ . Then  $\mathbf{X}'$  is the kernel of  $\widetilde{g}_X$  in  $C(\mathbf{P})$ . Hence

$$\text{pd Hom}_{\mathcal{D}}(\mathbf{P}, \mathbf{X})_B \leq \text{pd Hom}_{\mathcal{D}}(\mathbf{P}, \mathbf{X}')_B + 1.$$

Using induction on  $i$  and Lemma 3.2, we have that for  $\mathbf{X} \in C(\mathbf{P})$ , there is  $\mathbf{X}'$  such that

$$(1) \quad \mathbf{X}' \in C(\mathbf{P}) \cap (\text{add } H^0(\mathbf{P}) * \text{add } H^0(\mathbf{P})[1] * \cdots * \text{add } H^0(\mathbf{P})[d+1])$$

and  $\text{pd Hom}_{\mathcal{D}}(\mathbf{P}, \mathbf{X})_B \leq \text{pd Hom}_{\mathcal{D}}(\mathbf{P}, \mathbf{X}')_B + d + 1$ . By Lemma 2.4 and equation (1), we have  $\text{pd Hom}_{\mathcal{D}}(\mathbf{P}, \mathbf{X}')_B \leq d + 1$ . It then follows that  $\text{pd Hom}_{\mathcal{D}}(\mathbf{P}, \mathbf{X})_B \leq 2d + 2$ , and hence  $\text{gl.dim } B \leq 2d + 2$ .  $\square$

#### 4. THE CASE OF GLOBAL DIMENSION 2.

In this section, we consider the case when  $\text{gl.dim } A \leq 2$ . Our aim is to prove part (b) of Theorem 0.1, stating that in this case we have that the global dimension is at most 7 for the endomorphism algebra of any 2-term silting complex.

We prepare by showing four technical lemmas. Let  $\mathcal{P}_C^{[0,1]} = (\mathcal{P} * \mathcal{P}[1]) \cap C(\mathbf{P})$ .

**Lemma 4.1.** *If  $\mathbf{X}$  is in  $\mathcal{P}_C^{[0,1]}$ , then  $\text{pd Hom}_{\mathcal{D}}(\mathbf{P}, \mathbf{X})_B \leq 1$ .*

*Proof.* Since  $\mathbf{X}$  is in  $\mathcal{P} * \mathcal{P}[1]$ , there is a triangle  $\mathbf{O}_1 \rightarrow \mathbf{O}_0 \rightarrow \mathbf{X} \rightarrow \mathbf{O}_1[1]$  with  $\mathbf{O}_0, \mathbf{O}_1 \in \mathcal{P}$ . Applying the functor  $\text{Hom}_{\mathcal{D}}(\mathbf{P}, -)$  to this triangle, we get a long exact sequence

$$\text{Hom}_{\mathcal{D}}(\mathbf{P}, \mathbf{X}[-1]) \rightarrow \text{Hom}_{\mathcal{D}}(\mathbf{P}, \mathbf{O}_1) \rightarrow \text{Hom}_{\mathcal{D}}(\mathbf{P}, \mathbf{O}_0) \rightarrow \text{Hom}_{\mathcal{D}}(\mathbf{P}, \mathbf{X}) \rightarrow \text{Hom}_{\mathcal{D}}(\mathbf{P}, \mathbf{O}_1[1])$$

where the first term is zero since  $\mathbf{X}$  is in  $C(\mathbf{P})$ , and the last term is zero since  $\text{Hom}_{\mathcal{D}}(\mathbf{P}, \mathbf{P}[1]) = 0$ . Therefore,  $\text{pd Hom}_{\mathcal{D}}(\mathbf{P}, \mathbf{X})_B \leq 1$ .  $\square$

**Lemma 4.2.** *If  $\mathbf{X}$  is in  $C(\mathbf{P}) \cap (\mathcal{P}_C^{[0,1]} * \mathcal{P}_C^{[0,1]}[1] * \mathcal{P}_C^{[0,1]}[2])$ , then  $\text{pd Hom}_{\mathcal{D}}(\mathbf{P}, \mathbf{X})_B \leq 3$ .*

*Proof.* By  $\mathbf{X} \in \mathcal{P}_C^{[0,1]} * \mathcal{P}_C^{[0,1]}[1] * \mathcal{P}_C^{[0,1]}[2]$ , there are triangles

$$(2) \quad \mathbf{L} \rightarrow \mathbf{D}_1 \rightarrow \mathbf{X} \rightarrow \mathbf{L}[1]$$

and

$$(3) \quad \mathbf{D}_3 \rightarrow \mathbf{D}_2 \rightarrow \mathbf{L} \rightarrow \mathbf{D}_3[1]$$

with  $\mathbf{D}_1, \mathbf{D}_2, \mathbf{D}_3 \in \mathcal{P}_C^{[0,1]}$  and  $\mathbf{L} \in \mathcal{P}_C^{[0,1]} * \mathcal{P}_C^{[0,1]}[1] \subset \mathcal{P} * \mathcal{P}[1] * \mathcal{P}[2]$ . Applying  $\text{Hom}_{\mathcal{D}}(\mathbf{P}, -)$  to triangle (2), we obtain a long exact sequence

$$\begin{aligned} \text{Hom}_{\mathcal{D}}(\mathbf{P}, \mathbf{X}[-2]) &\rightarrow \text{Hom}_{\mathcal{D}}(\mathbf{P}, \mathbf{L}[-1]) \rightarrow \text{Hom}_{\mathcal{D}}(\mathbf{P}, \mathbf{D}_1[-1]) \rightarrow \text{Hom}_{\mathcal{D}}(\mathbf{P}, \mathbf{X}[-1]) \\ &\rightarrow \text{Hom}_{\mathcal{D}}(\mathbf{P}, \mathbf{L}) \rightarrow \text{Hom}_{\mathcal{D}}(\mathbf{P}, \mathbf{D}_1) \rightarrow \text{Hom}_{\mathcal{D}}(\mathbf{P}, \mathbf{X}) \rightarrow \text{Hom}_{\mathcal{D}}(\mathbf{P}, \mathbf{L}[1]). \end{aligned}$$

We have  $\text{Hom}_{\mathcal{D}}(\mathbf{P}, \mathbf{X}[i]) = 0$  for  $i = -1$  or  $i = -2$ , since  $\mathbf{X}$  is in  $C(\mathbf{P})$ . Furthermore, we have  $\text{Hom}_{\mathcal{D}}(\mathbf{P}, \mathbf{D}_1[-1]) = 0$ , by  $\mathbf{D}_1 \in C(\mathbf{P})$  and  $\text{Hom}_{\mathcal{D}}(\mathbf{P}, \mathbf{L}[1]) = 0$  by  $\mathbf{L} \in \mathcal{P} * \mathcal{P}[1] * \mathcal{P}[2]$ . From this it follows that we have a short exact sequence

$$0 \rightarrow \text{Hom}_{\mathcal{D}}(\mathbf{P}, \mathbf{L}) \rightarrow \text{Hom}_{\mathcal{D}}(\mathbf{P}, \mathbf{D}_1) \rightarrow \text{Hom}_{\mathcal{D}}(\mathbf{P}, \mathbf{X}) \rightarrow 0$$

and that  $\text{Hom}_{\mathcal{D}}(\mathbf{P}, \mathbf{L}[-1]) = 0$ . Using this short exact sequence, it follows that

$$(4) \quad \text{pd Hom}_{\mathcal{D}}(\mathbf{P}, \mathbf{X})_B \leq \max\{\text{pd Hom}_{\mathcal{D}}(\mathbf{P}, \mathbf{D}_1)_B, \text{pd Hom}_{\mathcal{D}}(\mathbf{P}, \mathbf{L})_B + 1\}.$$

Applying  $\text{Hom}_{\mathcal{D}}(\mathbf{P}, -)$  to the triangle (3), we obtain an exact sequence

$$0 = \text{Hom}_{\mathcal{D}}(\mathbf{P}, \mathbf{L}[-1]) \rightarrow \text{Hom}_{\mathcal{D}}(\mathbf{P}, \mathbf{D}_3) \rightarrow \text{Hom}_{\mathcal{D}}(\mathbf{P}, \mathbf{D}_2) \rightarrow \text{Hom}_{\mathcal{D}}(\mathbf{P}, \mathbf{L}) \rightarrow \text{Hom}_{\mathcal{D}}(\mathbf{P}, \mathbf{D}_3[1])$$

where the last term is zero due to  $\mathbf{D}_3 \in \mathcal{P}_C^{[0,1]}$ . As above, we obtain that

$$(5) \quad \text{pd Hom}_{\mathcal{D}}(\mathbf{P}, \mathbf{L})_B \leq \max\{\text{pd Hom}_{\mathcal{D}}(\mathbf{P}, \mathbf{D}_2)_B, \text{pd Hom}_{\mathcal{D}}(\mathbf{P}, \mathbf{D}_3)_B + 1\}.$$

Now, combining the inequalities (4) and (5) with Lemma 4.1, we obtain  $\text{pd Hom}_{\mathcal{D}}(\mathbf{P}, \mathbf{X})_B \leq 3$ .  $\square$

**Lemma 4.3.** *If  $\mathbf{N}$  is in  $\mathcal{D}^{\geq -1}(\mathbf{P}) \cap (\mathcal{P} * \mathcal{P}[1] * \mathcal{P}[2])$ , then there is an object  $\tilde{\mathbf{N}} \in C(\mathbf{P})$  such that  $\text{Hom}_{\mathcal{D}}(\mathbf{P}, \mathbf{N}) \cong \text{Hom}_{\mathcal{D}}(\mathbf{P}, \tilde{\mathbf{N}})$  as  $B$ -modules and  $\tilde{\mathbf{N}} \in \text{add } \mathbf{N} * \mathcal{P}_C^{[0,1]}[2]$ .*

*Proof.* Since  $(\mathcal{D}^{\leq 0}(\mathbf{P}), \mathcal{D}^{\geq 0}(\mathbf{P}))$  is a  $t$ -structure (see [12, Lemma 5.10]), there is a triangle

$$(6) \quad \mathbf{M} \rightarrow \mathbf{N} \rightarrow \tilde{\mathbf{N}} \rightarrow \mathbf{M}[1]$$

with  $\mathbf{M} \in \mathcal{D}^{\leq 0}(\mathbf{P})[1]$  and  $\tilde{\mathbf{N}} \in \mathcal{D}^{\geq 0}(\mathbf{P})$ . Then  $\mathbf{M} \in \mathcal{P}[1] * \dots * \mathcal{P}[l]$  for some  $l$  by [2, Proposition 2.23]. Applying the functor  $\text{Hom}_{\mathcal{D}}(\mathbf{P}, -)$  to the triangle (6), we have a long exact sequence

$$\dots \rightarrow \text{Hom}_{\mathcal{D}}(\mathbf{P}, \mathbf{M}[i]) \rightarrow \text{Hom}_{\mathcal{D}}(\mathbf{P}, \mathbf{N}[i]) \rightarrow \text{Hom}_{\mathcal{D}}(\mathbf{P}, \tilde{\mathbf{N}}[i]) \rightarrow \text{Hom}_{\mathcal{D}}(\mathbf{P}, \mathbf{M}[i+1]) \rightarrow \dots$$

Since  $\text{Hom}_{\mathcal{D}}(\mathbf{P}, \mathbf{M}[i]) = 0$  for  $i \geq 0$  by  $\mathbf{M} \in \mathcal{P}[1] * \dots * \mathcal{P}[l]$  and  $\text{Hom}_{\mathcal{D}}(\mathbf{P}, \tilde{\mathbf{N}}[i]) = 0$  for  $i < 0$  by  $\tilde{\mathbf{N}} \in \mathcal{D}^{\geq 0}(\mathbf{P})$ , and also  $\text{Hom}_{\mathcal{D}}(\mathbf{P}, \mathbf{N}[i]) = 0$  for  $i \neq -1, 0$  by the assumption  $\mathbf{N} \in \mathcal{D}^{\geq -1}(\mathbf{P}) \cap (\mathcal{P} * \mathcal{P}[1] * \mathcal{P}[2])$ , we have that

$$\text{Hom}_{\mathcal{D}}(\mathbf{P}, \mathbf{N}) \cong \text{Hom}_{\mathcal{D}}(\mathbf{P}, \tilde{\mathbf{N}})$$

as  $B$ -modules,

$$\text{Hom}_{\mathcal{D}}(\mathbf{P}, \tilde{\mathbf{N}}[i]) = 0, \text{ for } i > 0,$$

and

$$\text{Hom}_{\mathcal{D}}(\mathbf{P}, \mathbf{M}[i]) = 0, \text{ for } i < -1.$$

Thus, we obtain  $\tilde{\mathbf{N}} \in \mathcal{D}^{\leq 0}(\mathbf{P}) \cap \mathcal{D}^{\geq 0}(\mathbf{P}) = C(\mathbf{P})$  and  $\mathbf{M} \in \mathcal{D}^{\geq 0}(\mathbf{P})[1] \cap \mathcal{D}^{\leq 0}(\mathbf{P})[1] = C(\mathbf{P})[1]$ . Then by Lemma 2.3, we have  $\tilde{\mathbf{N}} \in \mathcal{P} * \mathcal{P}[1] * \mathcal{P}[2] * \mathcal{P}[3]$ . Applying the functor  $\text{Hom}_{\mathcal{D}}(-, \mathbf{P})$  to the triangle (6), we obtain a long exact sequence

$$\dots \rightarrow \text{Hom}_{\mathcal{D}}(\tilde{\mathbf{N}}, \mathbf{P}[i]) \rightarrow \text{Hom}_{\mathcal{D}}(\mathbf{N}, \mathbf{P}[i]) \rightarrow \text{Hom}_{\mathcal{D}}(\mathbf{M}, \mathbf{P}[i]) \rightarrow \text{Hom}_{\mathcal{D}}(\tilde{\mathbf{N}}, \mathbf{P}[i+1]) \rightarrow \dots$$

We have  $\text{Hom}_{\mathcal{D}}(\mathbf{N}, \mathbf{P}[i]) = 0$  for  $i > 2$  by  $\mathbf{N} \in \mathcal{P} * \mathcal{P}[1] * \mathcal{P}[2]$ , and we have  $\text{Hom}_{\mathcal{D}}(\tilde{\mathbf{N}}, \mathbf{P}[i]) = 0$  for  $i > 3$  by  $\tilde{\mathbf{N}} \in \mathcal{P} * \mathcal{P}[1] * \mathcal{P}[2] * \mathcal{P}[3]$ . From this it follows that  $\text{Hom}_{\mathcal{D}}(\mathbf{M}, \mathbf{P}[i]) = 0$  for  $i > 2$ , and hence  $\mathbf{M} \in (\mathcal{P}[1] * \mathcal{P}[2]) \cap C(\mathbf{P})[1] = \mathcal{P}_C^{[0,1]}[1]$ . Therefore we have that

$$\tilde{\mathbf{N}} \in \text{add } \mathbf{N} * \text{add } \mathbf{M}[1] \subset \text{add } \mathbf{N} * \mathcal{P}_C^{[0,1]}[2].$$

$\square$

**Lemma 4.4.** *Let  $\mathbf{X} \in C(\mathbf{P}) \cap (\mathcal{P} * \mathcal{P}[1] * \cdots * \mathcal{P}[t] * \mathcal{H}[t+1])$  for some  $t$  with  $0 \leq t \leq 3$ , where  $\mathcal{H} \subset \mathcal{P} * \mathcal{P}[1] * \cdots * \mathcal{P}[2-t]$  (and  $\mathcal{H} = 0$  for  $t = 3$ ). Then for each  $r$  with  $0 \leq r \leq \min\{t+1, 3\}$ , there is an object  $\widetilde{\mathbf{X}}_r \in C(\mathbf{P})$  such that*

$$\mathrm{pd} \mathrm{Hom}_{\mathcal{D}}(\mathbf{P}, \mathbf{X})_B \leq \mathrm{pd} \mathrm{Hom}_{\mathcal{D}}(\mathbf{P}, \widetilde{\mathbf{X}}_r)_B + r$$

and

$$\widetilde{\mathbf{X}}_r \in \mathcal{P} * \mathcal{P}[1] * \cdots * \mathcal{P}[t-r] * \mathcal{H}[t+1-r] * \mathcal{P}_C^{[0,1]}[3-r] * \cdots * \mathcal{P}_C^{[0,1]}[2]$$

where  $\mathcal{P} * \mathcal{P}[1] * \cdots * \mathcal{P}[t-r]$  is taken to be 0 when  $r = t+1$  and  $\mathcal{P}_C^{[0,1]}[3-r] * \cdots * \mathcal{P}_C^{[0,1]}[2]$  is taken to be 0 when  $r = 0$ .

*Proof.* Let  $\widetilde{\mathbf{X}}_0 = \mathbf{X}$ . Then  $\widetilde{\mathbf{X}}_0$  satisfies the conditions in the lemma. Assume that  $\widetilde{\mathbf{X}}_{r-1}$  satisfying the conditions. By

$$\widetilde{\mathbf{X}}_{r-1} \in \mathcal{P} * \mathcal{P}[1] * \cdots * \mathcal{P}[t-(r-1)] * \mathcal{H}[t+1-(r-1)] * \mathcal{P}_C^{[0,1]}[3-(r-1)] * \cdots * \mathcal{P}_C^{[0,1]}[2],$$

there is a triangle

$$\mathbf{X}_r \rightarrow \mathbf{P}_0 \rightarrow \widetilde{\mathbf{X}}_{r-1} \rightarrow \mathbf{X}_r[1]$$

with  $\mathbf{P}_0 \in \mathcal{P}$ ,  $\mathbf{X}_r \in \mathcal{P} * \cdots * \mathcal{P}[t-r] * \mathcal{H}[t+1-r] * \mathcal{P}_C^{[0,1]}[3-r] * \cdots * \mathcal{P}_C^{[0,1]}[1] \subset \mathcal{P} * \mathcal{P}[1] * \mathcal{P}[2]$ . The inclusion follows from Lemma 1.2. Applying  $\mathrm{Hom}_{\mathcal{D}}(\mathbf{P}, -)$  to this triangle, we have a long exact sequence

$$\cdots \rightarrow \mathrm{Hom}_{\mathcal{D}}(\mathbf{P}, \mathbf{X}_r[i]) \rightarrow \mathrm{Hom}_{\mathcal{D}}(\mathbf{P}, \mathbf{P}_0[i]) \rightarrow \mathrm{Hom}_{\mathcal{D}}(\mathbf{P}, \widetilde{\mathbf{X}}_{r-1}[i]) \rightarrow \mathrm{Hom}_{\mathcal{D}}(\mathbf{P}, \mathbf{X}_r[i+1]) \rightarrow \cdots$$

Since  $\mathrm{Hom}_{\mathcal{D}}(\mathbf{P}, \widetilde{\mathbf{X}}_{r-1}[i]) = 0$  for  $i \neq 0$  by  $\widetilde{\mathbf{X}}_{r-1} \in C(\mathbf{P})$ ,  $\mathrm{Hom}_{\mathcal{D}}(\mathbf{P}, \mathbf{X}_r[1]) = 0$  by  $\mathbf{X}_r \in \mathcal{P} * \mathcal{P}[1] * \mathcal{P}[2]$ , and also  $\mathrm{Hom}_{\mathcal{D}}(\mathbf{P}, \mathbf{P}_0[i]) = 0$  for  $i < -1$  by  $\mathbf{P}$  being 2-term, we have a short exact sequence

$$0 \rightarrow \mathrm{Hom}_{\mathcal{D}}(\mathbf{P}, \mathbf{X}_r) \rightarrow \mathrm{Hom}_{\mathcal{D}}(\mathbf{P}, \mathbf{P}_0) \rightarrow \mathrm{Hom}_{\mathcal{D}}(\mathbf{P}, \widetilde{\mathbf{X}}_{r-1}) \rightarrow 0$$

and

$$\mathrm{Hom}_{\mathcal{D}}(\mathbf{P}, \mathbf{X}_r[i]) = 0 \text{ for } i < -1.$$

Then  $\mathrm{pd} \mathrm{Hom}_{\mathcal{D}}(\mathbf{P}, \widetilde{\mathbf{X}}_{r-1})_B \leq \mathrm{pd} \mathrm{Hom}_{\mathcal{D}}(\mathbf{P}, \mathbf{X}_r)_B + 1$  and by Lemma 4.3, there is an object  $\widetilde{\mathbf{X}}_r \in C(\mathbf{P})$  such that  $\mathrm{Hom}_{\mathcal{D}}(\mathbf{P}, \widetilde{\mathbf{X}}_r)_B \cong \mathrm{Hom}_{\mathcal{D}}(\mathbf{P}, \mathbf{X}_r)_B$  and

$$\widetilde{\mathbf{X}}_r \in \mathrm{add} \mathbf{X}_r * \mathcal{P}_C^{[0,1]}[2] \subset \mathcal{P} * \mathcal{P}[1] * \cdots * \mathcal{P}[t-r] * \mathcal{H}[t+1-r] * \mathcal{P}_C^{[0,1]}[3-r] * \cdots * \mathcal{P}_C^{[0,1]}[1] * \mathcal{P}_C^{[0,1]}[2].$$

□

Now we prove the main result in this section.

**Theorem 4.5.** *If  $\mathrm{gl.dim} A \leq 2$ , then  $\mathrm{gl.dim} \mathrm{End}_{\mathcal{D}}(\mathbf{P}) \leq 7$  for any 2-term silting complex  $\mathbf{P}$  in  $K^b(\mathrm{proj} A)$ .*

*Proof.* Let  $\mathbf{X} \in C(\mathbf{P})$ . Then by Lemma 2.3, we have that  $\mathbf{X} \in \mathcal{P} * \mathcal{P}[1] * \mathcal{P}[2] * \mathcal{P}[3]$ . By Lemma 4.4, (taking  $t = 3$ ,  $r = 2$  and hence  $\mathcal{H} = 0$ ), there is an  $\widetilde{\mathbf{X}} \in C(\mathbf{P})$  such that  $\widetilde{\mathbf{X}} \in \mathcal{P} * \mathcal{P}[1] * \mathcal{P}_C^{[0,1]}[1] * \mathcal{P}_C^{[0,1]}[2]$ , and

$$(7) \quad \mathrm{pd} \mathrm{Hom}_{\mathcal{D}}(\mathbf{P}, \mathbf{X})_B \leq \mathrm{pd} \mathrm{Hom}_{\mathcal{D}}(\mathbf{P}, \widetilde{\mathbf{X}})_B + 2.$$

Then there is a triangle

$$\mathbf{Z} \rightarrow \mathbf{Y} \rightarrow \widetilde{\mathbf{X}} \rightarrow \mathbf{Z}[1]$$

with  $\mathbf{Y} \in \mathcal{P} * \mathcal{P}[1]$  and  $\mathbf{Z} \in \mathcal{P}_C^{[0,1]} * \mathcal{P}_C^{[0,1]}[1]$ . Applying the functor  $\mathrm{Hom}_{\mathcal{D}}(\mathbf{P}, -)$  to this triangle, we have a long exact sequence

$$\cdots \rightarrow \mathrm{Hom}_{\mathcal{D}}(\mathbf{P}, \mathbf{Z}[i]) \rightarrow \mathrm{Hom}_{\mathcal{D}}(\mathbf{P}, \mathbf{Y}[i]) \rightarrow \mathrm{Hom}_{\mathcal{D}}(\mathbf{P}, \widetilde{\mathbf{X}}[i]) \rightarrow \mathrm{Hom}_{\mathcal{D}}(\mathbf{P}, \mathbf{Z}[i+1]) \rightarrow \cdots$$

Since  $\text{Hom}_{\mathcal{D}}(\mathbf{P}, \widetilde{\mathbf{X}}[i]) = 0$  for  $i \neq 0$  by  $\widetilde{\mathbf{X}} \in C(\mathbf{P})$ , and  $\text{Hom}_{\mathcal{D}}(\mathbf{P}, \mathbf{Z}[i]) = 0$  for  $i \neq -1, 0$  by  $\mathbf{Z} \in C(\mathbf{P}) * C(\mathbf{P})[1]$ , we have a short exact sequence

$$0 \rightarrow \text{Hom}_{\mathcal{D}}(\mathbf{P}, \mathbf{Z}) \rightarrow \text{Hom}_{\mathcal{D}}(\mathbf{P}, \mathbf{Y}) \rightarrow \text{Hom}_{\mathcal{D}}(\mathbf{P}, \widetilde{\mathbf{X}}) \rightarrow 0,$$

and

$$\text{Hom}_{\mathcal{D}}(\mathbf{P}, \mathbf{Y}[i]) = 0 \text{ for } i < -1.$$

Then we have that

$$(8) \quad \text{pd Hom}_{\mathcal{D}}(\mathbf{P}, \widetilde{\mathbf{X}})_B \leq \max\{\text{pd Hom}_{\mathcal{D}}(\mathbf{P}, \mathbf{Y})_B, \text{pd Hom}_{\mathcal{D}}(\mathbf{P}, \mathbf{Z})_B + 1\}$$

and  $\mathbf{Y}, \mathbf{Z} \in \mathcal{D}^{\geq -1}(\mathbf{P})$ . By Lemma 4.3, there are objects  $\widetilde{\mathbf{Y}}, \widetilde{\mathbf{Z}} \in C(\mathbf{P})$  such that:

$$\widetilde{\mathbf{Y}} \in \mathcal{P} * \mathcal{P}[1] * \mathcal{P}_C^{[0,1]}[2] \quad \widetilde{\mathbf{Z}} \in \mathcal{P}_C^{[0,1]} * \mathcal{P}_C^{[0,1]}[1] * \mathcal{P}_C^{[0,1]}[2]$$

$$\text{Hom}_{\mathcal{D}}(\mathbf{P}, \widetilde{\mathbf{Y}}_B) \cong \text{Hom}_{\mathcal{D}}(\mathbf{P}, \mathbf{Y})_B \quad \text{Hom}_{\mathcal{D}}(\mathbf{P}, \widetilde{\mathbf{Z}}_B) \cong \text{Hom}_{\mathcal{D}}(\mathbf{P}, \mathbf{Z})_B$$

By Lemma 4.4, (taking  $t = 1, r = 2$  and  $\mathcal{H} = \mathcal{P}_C^{[0,1]}[2]$ ), there is an object  $\widetilde{\mathbf{Y}}' \in C(\mathbf{P})$  such that  $\widetilde{\mathbf{Y}}' \in \mathcal{P}_C^{[0,1]} * \mathcal{P}_C^{[0,1]}[1] * \mathcal{P}_C^{[0,1]}[2]$  and

$$(9) \quad \text{pd Hom}_{\mathcal{D}}(\mathbf{P}, \widetilde{\mathbf{Y}})_B \leq \text{pd Hom}_{\mathcal{D}}(\mathbf{P}, \widetilde{\mathbf{Y}}')_B + 2.$$

By Lemma 4.2, we have  $\text{pd Hom}_{\mathcal{D}}(\mathbf{P}, \widetilde{\mathbf{Z}})_B \leq 3$  and  $\text{pd Hom}_{\mathcal{D}}(\mathbf{P}, \widetilde{\mathbf{Y}}')_B \leq 3$ . Hence, combining (7), (8) and (9), we obtain

$$\begin{aligned} \text{pd Hom}_{\mathcal{D}}(\mathbf{P}, \mathbf{X})_B &\leq \text{pd Hom}_{\mathcal{D}}(\mathbf{P}, \widetilde{\mathbf{X}})_B + 2 \\ &\leq \max\{\text{pd Hom}_{\mathcal{D}}(\mathbf{P}, \mathbf{Y})_B, \text{pd Hom}_{\mathcal{D}}(\mathbf{P}, \mathbf{Z})_B + 1\} + 2 \\ &= \max\{\text{pd Hom}_{\mathcal{D}}(\mathbf{P}, \widetilde{\mathbf{Y}})_B, \text{pd Hom}_{\mathcal{D}}(\mathbf{P}, \widetilde{\mathbf{Z}})_B + 1\} + 2 \\ &\leq \max\{\text{pd Hom}_{\mathcal{D}}(\mathbf{P}, \widetilde{\mathbf{Y}}')_B + 2, \text{pd Hom}_{\mathcal{D}}(\mathbf{P}, \widetilde{\mathbf{Z}})_B + 1\} + 2 \\ &\leq 7. \end{aligned}$$

□

## 5. EXAMPLES

**5.1. First example.** We first give an example to show that the bound in Theorem 3.3 is possible. Let  $n \geq 2$  and  $A = \mathbf{k}Q/I$  where  $Q$  is the following quiver

$$\begin{array}{ccccccc} 2n+3 & \xrightarrow{a_n} & \cdots & \xrightarrow{a_3} & 7 & \xrightarrow{a_2} & 5 \xrightarrow{a_1} 3 \\ & & & & & & \searrow c_1 \\ & & & & & & 2 \xrightarrow{c_2} 1 \\ & & & & & & \nearrow d_2 \\ & & & & & & 4 \xrightarrow{d_1} 1 \\ 2n+4 & \xleftarrow{b_n} & \cdots & \xleftarrow{b_3} & 8 & \xleftarrow{b_2} & 6 \xleftarrow{b_1} 4 \end{array}$$

and the ideal  $I$  is generated by  $c_1d_1 - c_2d_2$ ,  $a_{i+1}a_i$  and  $b_i b_{i+1}$ ,  $1 \leq i \leq n-1$ . Then  $\text{gl.dim } A = n$ . Let  $\mathbf{P}$  be the direct sum of the following complexes in  $K^b(\text{proj } A)$ :

$$\begin{array}{ccc} 0 & \longrightarrow & \bigoplus_{1 \leq i \leq n+2, i \neq 2} P_{2i} \\ P_4 & \longrightarrow & P_2 \\ P_1 & \longrightarrow & P_3 \\ \bigoplus_{1 \leq i \leq n+2, i \neq 2} P_{2i-1} & \longrightarrow & 0 \end{array} \quad ,$$

It is easily verified that  $\mathbf{P}$  is a 2-term silting complex. The quiver of the endomorphism ring  $\text{End}_{\mathcal{D}}(\mathbf{P})$  is the Dynkin quiver of type  $A_{2n+4}$ :

$$\begin{array}{ccccccccccc} 2n+3 & \xrightarrow{a_n} & \cdots & \xrightarrow{a_3} & 7 & \xrightarrow{a_2} & 5 & \xrightarrow{a_1} & 3 & \xrightarrow{a_0} & 1 \\ & & & & & & & & & & \downarrow c \\ 2n+4 & \xleftarrow{b_n} & \cdots & \xleftarrow{b_3} & 8 & \xleftarrow{b_2} & 6 & \xleftarrow{b_1} & 4 & \xleftarrow{b_0} & 2 \end{array}$$

with the relations  $a_{i+1}a_i = 0$ ,  $b_i b_{i+1} = 0$ ,  $1 \leq i \leq n-1$  and  $a_0 c b_0 = 0$ . Hence the global dimension of  $\text{End}_{\mathcal{D}}(\mathbf{P})$  is  $2n+2$ .

**5.2. Second example.** The next example shows that 7 is a possible value for the global dimension of the endomorphism algebra of a 2-term silting complex over an algebra of global dimension two. Let  $A = \mathbf{k}Q/I$  with  $Q$  the following quiver

$$1 \xrightarrow{a_1} 2 \xrightarrow{a_2} 3 \xrightarrow{a_3} 4 \xrightleftharpoons[a_5]{a_4} 5 \xrightarrow{a_6} 6 \xrightarrow{a_7} 7 \xrightarrow{a_8} 8$$

and  $I$  the ideal generated by  $a_1 a_2$ ,  $a_3 a_4 a_6$  and  $a_7 a_8$ . Then  $A$  has global dimension two, and the complex  $\mathbf{P}$  given by the direct sum of the complexes

$$\begin{array}{ccc} 0 & \longrightarrow & \bigoplus_{i=5,7,8} P_i \\ P_6 & \longrightarrow & P_5 \\ P_4 & \longrightarrow & P_3 \\ \bigoplus_{i=1,2,4} P_i & \longrightarrow & 0 \end{array}.$$

is a 2-term silting complex. It is easily verified, that the quiver of  $\text{End}_{\mathcal{D}}(\mathbf{P})$  is a linearly oriented Dynkin quiver of type  $A_8$  and the ideal of relation equals the square of the Jacobson radical. Hence the global dimension of  $\text{End}_{\mathcal{D}}(\mathbf{P})$  is 7.

**5.3. Third example.** The last example shows that there is no bound on the global dimension of the endomorphism algebra of a 2-silting object over an algebra with global dimension  $d \geq 3$ . This example then completes the proof of Theorem 0.1.

Let first  $A = \mathbf{k}Q/I$  where the quiver  $Q$  is

$$\begin{array}{ccccc} 3 & \xleftarrow{a} & 2 & \xleftarrow{e} & 4 \\ & \xrightarrow{b} & & \xrightarrow{d} & \\ & & \downarrow c & & \\ & & 1 & & \end{array}$$

and  $I = \langle ba, bd, abc, de \rangle$ . The indecomposable projective  $A$ -modules are

$$P_1 = 1, \quad P_2 = \begin{smallmatrix} 2 \\ 1 \end{smallmatrix} \begin{smallmatrix} 3 \\ 2 \end{smallmatrix} 4, \quad P_3 = \begin{smallmatrix} 3 \\ 2 \end{smallmatrix}, \quad P_4 = \begin{smallmatrix} 4 \\ 1 \end{smallmatrix} \begin{smallmatrix} 2 \\ 3 \end{smallmatrix} 4.$$

The integers here denote the corresponding simples, and the notation indicates the radical filtration. The global dimension of  $A$  is 3. Let  $\mathbf{P}$  be the direct sum of

$$\mathbf{P}_i = \cdots \rightarrow 0 \rightarrow P_i \rightarrow 0 \rightarrow 0 \rightarrow \cdots, \quad i = 1, 3, 4,$$

(concentrated in degree -1) and

$$\mathbf{P}_2 = \cdots \rightarrow 0 \rightarrow P_1 \oplus P_3 \oplus P_4 \xrightarrow{p} P_2 \rightarrow 0 \rightarrow \cdots$$

where  $p$  is a projective presentation of  $S_2$ . Then it is easily verified that  $\mathbf{P}$  is a 2-term silting complex.

By Proposition 1.1(b) we have that  $\mathcal{T}(\mathbf{P}) = \text{Fac } H^0(\mathbf{P})$ , and hence  $\mathcal{T}(\mathbf{P}) = \text{add } S_2$ . We will show the projective dimension of  $S_2$  in  $C(\mathbf{P})$  is infinite, by proving that its third syzygy equals  $S_2$ . This implies that a minimal projective resolution of  $S_2$  is periodic and hence infinite.

Using the notation in Section 2, we have that  $\tilde{\mathbf{P}}_1 = \mathbf{P}_1$  and  $\tilde{\mathbf{P}}_3 = \mathbf{P}_3$ . Moreover  $\tilde{\mathbf{P}}_4 = (P_4/S_2)[1]$ , and  $\tilde{\mathbf{P}}_2$  is given by the complex

$$\cdots \rightarrow 0 \rightarrow P_1 \oplus P_3 \oplus (P_4/S_2) \xrightarrow{\tilde{p}} P_2 \rightarrow 0 \rightarrow \cdots$$

Consider now the triangle

$$\text{cone}(\pi)[-1] \rightarrow \tilde{\mathbf{P}}_2 \xrightarrow{\pi} S_2 \rightarrow \text{cone}(\pi)$$

where  $\pi$  is

$$\begin{array}{ccccccc} \cdots & \rightarrow & 0 & \rightarrow & P_1 \oplus P_3 \oplus (P_4/S_2) & \xrightarrow{\tilde{p}} & P_2 \rightarrow 0 \rightarrow \cdots \\ & & \downarrow & & \downarrow & & \pi^0 \downarrow \quad \downarrow \\ \cdots & \rightarrow & 0 & \rightarrow & 0 & \rightarrow & S_2 \rightarrow 0 \rightarrow \cdots \end{array}$$

with  $\pi^0$  being a projective cover of  $S_2$  in  $\text{mod } A$ .

Then  $H^0(\text{cone}(\pi)[-1]) = 0$ ,  $H^{-1}(\text{cone}(\pi)[-1]) \cong H^{-1}(\tilde{\mathbf{P}}_2) \in \mathcal{F}(\mathbf{P})$  and  $H^i(\text{cone}(\pi)[-1]) = 0$  for  $i \neq -1, 0$ . So  $\text{cone}(\pi)[-1]$  is in  $C(\mathbf{P})$ , using Proposition 1.1 (d). Hence  $\pi$  is a projective cover of  $S_2$  in  $C(\mathbf{P})$  and  $\text{cone}(\pi)[-1]$  is its kernel in  $C(\mathbf{P})$ .

Note that  $\text{cone}(\pi)[-1] \cong H^{-1}(\text{cone}(\pi)[-1]) \cong H^{-1}(\tilde{\mathbf{P}}_2) \cong P_1 \oplus M$ , where  $M = {}_1^2_3 4 \in \mathcal{F}(\mathbf{P})$ . Consider the triangle

$$\text{cone}(\pi_1)[-1] \rightarrow \tilde{\mathbf{P}}_1 \oplus \tilde{\mathbf{P}}_3 \oplus \tilde{\mathbf{P}}_4 \xrightarrow{\pi_1} M[1] \rightarrow \text{cone}(\pi_1)$$

where  $\pi_1$  is

$$\begin{array}{ccccccc} \cdots & \rightarrow & 0 & \rightarrow & P_1 \oplus P_3 \oplus (P_4/S_2) & \rightarrow & 0 \rightarrow 0 \rightarrow \cdots \\ & & \downarrow & & \pi_1^{-1} \downarrow & & \downarrow \quad \downarrow \\ \cdots & \rightarrow & 0 & \rightarrow & M & \rightarrow & 0 \rightarrow 0 \rightarrow \cdots \end{array}$$

with  $\pi_1^{-1}$  being the unique (up to a scalar) right minimal homomorphism from  $P_1 \oplus P_3 \oplus (P_4/S_2)$  to  $M$ . Then  $H^0(\text{cone}(\pi_1)[-1]) \cong S_2 \in \mathcal{T}(\mathbf{P})$ ,  $H^{-1}(\text{cone}(\pi_1)[-1]) \cong {}_1^2 \oplus M \in \mathcal{F}(\mathbf{P})$  (since  $\text{Hom}_A(S_2, {}_1^2) = 0$ ) and  $H^i(\text{cone}(\pi_1)[-1]) = 0$  for  $i \neq -1, 0$ . So  $\text{cone}(\pi_1)[-1] \in C(\mathbf{P})$ , hence  $\pi_1$  is a projective cover of  $M[1]$  in  $C(\mathbf{P})$  and  $\text{cone}(\pi_1)[-1]$  is its kernel.

Now consider the triangle

$$\text{cone}(\pi_2)[-1] \rightarrow \tilde{\mathbf{P}}_2 \xrightarrow{\pi_2} \text{cone}(\pi_1)[-1] \rightarrow \text{cone}(\pi_2)$$

where  $\pi_2$  is

$$\begin{array}{ccccccc} \cdots & \rightarrow & 0 & \rightarrow & P_1 \oplus P_3 \oplus (P_4/S_2) & \xrightarrow{p} & P_2 \rightarrow 0 \rightarrow \cdots \\ & & \downarrow & & \pi_2^{-1} \downarrow & & \pi_2^0 \downarrow \quad \downarrow \\ \cdots & \rightarrow & 0 & \rightarrow & P_1 \oplus P_3 \oplus (P_4/S_2) & \xrightarrow{\pi_1^{-1}} & M \rightarrow 0 \rightarrow \cdots \end{array}$$

with  $\pi_2^{-1}$  being the identity map and  $\pi_2^0$  being a projective cover of  $M$  in  $\text{mod } A$ . Then we have  $H^0(\text{cone}(\pi_2)[-1]) \cong S_2 \in \mathcal{T}(\mathbf{P})$  and  $H^i(\text{cone}(\pi_2)[-1]) = 0$  for  $i \neq 0$ . Hence  $\text{cone}(\pi_2)[-1] \cong S_2 \in C(\mathbf{P})$  and we have a short exact sequence in  $C(\mathbf{P})$ :

$$0 \rightarrow S_2 \rightarrow \tilde{\mathbf{P}}_2 \xrightarrow{\pi_2} (\text{cone}(\pi_1)[-1]) \rightarrow 0.$$

Thus, the projective resolution of  $S_2$  in  $C(\mathbf{P})$  is periodic and hence the projective dimension is infinite. Therefore, also the global dimension of  $B$  is infinite, by Proposition 1.1 (e).

Now, for any  $n$  consider the quiver  $Q_n$  given by

$$\begin{array}{ccccccc}
 3 & \xrightleftharpoons[b]{a} & 2 & \xrightleftharpoons[d]{e} & 4 & & \\
 & & \downarrow c_0 & & & & \\
 & & 1_0 & \xrightarrow{c_1} & 1_1 & \xrightarrow{c_2} & 1_2 \xrightarrow{c_3} \cdots \xrightarrow{c_n} 1_n
 \end{array}$$

with relations  $I_n = \langle ba, bd, abc_0, de, c_0c_1, c_1c_2, \dots, c_{n-1}c_n \rangle$ . Consider the algebra  $A(n) = kQ_n/I_n$ . We leave it as an exercise to check that  $A(n)$  has global dimension  $n + 3$ , and to find a 2-term silting complex  $\mathbf{P}'$ , such that  $\text{End}_{D^b(A(n))}(\mathbf{P}')$  has infinite global dimension.

#### REFERENCES

- [1] T. Adachi, O. Iyama and I. Reiten,  $\tau$ -tilting theory, *Compos. Math.* 150 (2015), no. 3, 415–452.
- [2] T. Aihara and O. Iyama, *Silting mutation in triangulated categories*, *J. Lond. Math. Soc.* (2) 85 (2012), no. 3, 633–668.
- [3] I. Assem, D. Simson and A. Skowronski, *Elements of the representation theory of associative algebras. Vol. 1. Techniques of Representation Theory*, London Mathematical Society Student Texts, 65. Cambridge University Press, Cambridge, 2006.
- [4] T. Brüstle and D. Yang, *Ordered Exchange Graphs*, *Advances in representation theory of algebras*, 135–193, EMS Ser. Congr. Rep., Eur. Math. Soc., Zürich, 2013.
- [5] A. B. Buan and Y. Zhou, *A silting theorem*, *J. Pure Appl. Algebra* 220 (2016), no. 7, 2748–2770.
- [6] A. B. Buan and Y. Zhou, *Silted algebras*, preprint arXiv:1506.03649.
- [7] F. U. Coelho and M. A. Lanzilotta, *Algebras with small homological dimensions*, *Manuscripta Mathematica* 100 (1999) 1–11.
- [8] D. Happel, *Triangulated categories in the representation theory of finite dimensional algebras*, London Mathematical Society Lecture Note Series, 119, Cambridge University Press, Cambridge, 1988.
- [9] D. Happel, I. Reiten and S. O. Smalø, *Tilting in abelian categories and quasitilted algebra*, *Mem. Amer. Math. Soc.* 120 (1996), no. 575.
- [10] O. Iyama and Y. Yoshino, *Mutation in triangulated categories and rigid Cohen-Macaulay modules*, *Invent. Math.* 172 (2008), no. 1, 117–168.
- [11] B. Keller and D. Vossieck, *Aisles in derived categories*, *Bull. Soc. Math. Belg. Sr. A* 40 (1988), no. 2, 239–253.
- [12] S. Koenig and D. Yang, *Silting objects, simple-minded collections, t-structures and co-t-structures for finite-dimensional algebras*, *Doc. Math.* 19 (2014), 403–438.
- [13] J. Rickard, *Morita theory for derived categories*, *J. London Math. Soc.* (2) 39 (1989), no. 3, 436–456.

DEPARTMENT OF MATHEMATICAL SCIENCES NORWEGIAN UNIVERSITY OF SCIENCE AND TECHNOLOGY 7491 TRONDHEIM NORWAY

*E-mail address:* aslabb@math.ntnu.no

DEPARTMENT OF MATHEMATICAL SCIENCES NORWEGIAN UNIVERSITY OF SCIENCE AND TECHNOLOGY 7491 TRONDHEIM NORWAY

*E-mail address:* yu.zhou@math.ntnu.no